



Profiles in Error: A Template for Quality Improvement

Portelli K¹, Farrell A¹, Dechario SP², Rindskopf D³, Husk G¹, Ansari P¹, Andrews R¹, Carroccio A¹, Giangola G¹, Antonacci AC^{4,*}

¹Lenox Hill Hospital, Department of Surgery, USA

²Institute for Spine and Scoliosis, USA

³Graduate School, City University of New York, USA

⁴Weill Cornell Medical College, USA

***Correspondence:**

Anthony C Antonacci, MD, SM, FACS
Weill Cornell Medical College, 1300 York Avenue, New York, NY 10065, USA.
Email: aantonacci@gmail.com

Article Info:

Received Date: May 04, 2026

Published Date: May 25, 2026

Citation:

Portelli K, Farrell A, Dechario SP, Rindskopf D, Husk G, Ansari P, Andrews R, Carroccio A, Giangola G, Antonacci AC. Profiles in Error: A Template for Quality Improvement. J Surg Operative Tech. 2026;1(1):1-6.

Copyright © Anthony C Antonacci

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.



ABSTRACT

Introduction: It has been alleged that human error is impossible to eliminate in surgery because of the complex, multifactorial nature of clinical care. However, the level of detail that describes error types is limited and studies analyzing large numbers of cases are few. We developed a standardized error template and gathered data over five years to profile patterns of error in surgical cases with complications.

Methods: Data was collected at an accredited general surgery residency program utilizing a previously described complication reporting system. The system allows for click-based, electronic entry of complications, case summaries, and a standardized critique algorithm across multiple domains including diagnostic, technical, judgment, therapy/prophylaxis, communication, professionalism and system error. Data was collected on cases with complications and reviewed at weekly validation meetings by a committee of attendings (including the site Chief of Surgery, Program Director, divisional leadership, senior and junior residents, nursing and quality). The process was prospective and comprehensive; presented by residents responsible for care; designed to validate complications and identify error. Data was analyzed for the incidence of error across the seven domains, thirty-seven subtypes, and for specific procedures including bariatrics, cholecystectomy, colectomy, hepatic procedures, hiatal hernia, ileostomy closure, pancreatectomy and ventral hernia/component cases.

Results: The study population included 25,857 general surgery/vascular cases, 4,005 complications, 1,473 cases with complications, and 3,268 errors or 2.2 errors per complication case. The number and incidence of error by domain category was technical (831, 25.4%), therapy/prophylaxis (598, 18.3%), judgment (509, 15.6%), diagnosis (378, 11.6%), system issues (379, 11.3%), communication (356, 10.9%), and professionalism (226, 6.9%). The most common subtypes for each domain were ‘error technical performance’(54.2%), ‘post procedure care MD Team’(79.4%), ‘error medical judgment’(50.5%), ‘delay diagnosis’(11.6%), ‘pre/post procedure care Nursing Team’(59.7%), ‘error in handoff team members’(10.9%), and ‘identified/accepted

responsibility'(62%), respectively. All of the operative procedures evaluated mimicked the error profile of the domain and subtype categories with some differences in emphasis. We calculated the proportion of operations with complications that had errors. The median sub-specialty had a rate of .83, with a range of only .77 to .89. Thus, a large proportion of operations with complications had at least one error, and this did not vary much across sub-specialty.

Conclusion: These data suggest that patterns of error can be accurately described utilizing a standardized error profile template to analyze post-operative complications. Operation types with more errors had more complications. By understanding the distribution of error types, providers and colleagues may be informed of their individual propensity for error, and systems can be more focused in quality improvement initiatives based on patterns specific to procedure type.

KEYWORDS: Quality Improvement (QI), Error, Harm

INTRODUCTION

Medical error is reported to be the third leading cause of death in the United States, with estimates suggesting it accounts for a significant portion of preventable morbidity and mortality.¹ Despite prominence of this statistic, the definition of "error" remains a widely debated topic in the healthcare community. Probably the most famous definition of error was set out in the landmark publication "To Err is Human" 1999 which asserts that error is "either a failure of a planned action to be completed as intended, or the use of a wrong plan to achieve an aim."^{2,3} In other words, error is a result of the human condition and inevitable in any field wherein human reason and decision-making impact outcomes. Based on this definition, it has been accepted that there will always be some component of human error in medicine.

Further studies have worked to define error into more quantifiable measures, creating classification and categorization systems to stratify and study different types of error with the aim of accurately describing and minimizing error, thereby reducing patient complications. Some systems are intention based while others remain more objective. However, error reporting remains inconsistent, and

few large-scale trials exist. In addition, while it should be evaluated, not all errors lead to harm. Unlike medical error, which encompasses all mistakes made in the field, harm is defined as the error that reaches the patient and results in a negative or adverse outcome.

More recently, researchers have been interested in determining the root causes of medical error. In many studies, error has been classified by pre-determined categories. For instance, Marsh et al performed a systematic review of the literature related to medical mistakes, finding a myriad of disparate definitions of medical error. At the conclusion of the review, the authors proposed five standardized categories of error including: medication errors, errors in diagnosis, technical errors, system errors, and judgment errors.⁴ Moreover, studies of error have sought to relate errors to specific procedures, recognizing that different surgeries are prone to particular errors and may not be easily compared. Using both the aforementioned categorization system and surgical subgroup stratification, subsequent research efforts by Marsh focused on role of judgement errors in harmful surgical outcomes from medical mistakes.¹

In this paper, we present a set standardized error template and evaluate data gathered over five years to profile patterns of error in surgical cases with complications. We hypothesized that errors leading to harm are not typically attributable to a single source but rather arise from complex, multifactorial sources including human, systemic and procedural elements.

METHODS

Data was collected at an accredited general surgery residency program utilizing a complication reporting system previously described.⁵⁻¹¹ The system allows for web-based electronic data entry system and collects complication data via clinical case summaries and a 90% click-based data entry system. This allows for standardized identification of over 300 surgical complications (Figure 1). The system requires attending, team and patient identifiers. It begins with a detailed case summary of the case events followed by a series of click-based inputs which includes a standardized critique algorithm for error. The error domains included: diagnostic, technical, judgment, therapy/prophylaxis, communication, professionalism and system error (Figure 2).

The screenshot shows a web-based interface for reporting complications. At the top, there's a header 'Attending Evaluation: Resident Critique'. Below it are several input fields: Surgeon (with a dropdown), DOB, AGE, SEX, MRN, DOS, Procedure, Run Time, Srg Time, and Case Type. To the right, there's a section for 'Currently Selected Complications:' with a dropdown menu. Below these fields are four buttons: 'Delete Selected' (blue), 'Email Resident Grade' (green), 'Save And Back' (red), and 'Cancel' (yellow). The main content area is a grid of five columns, each representing a different category of complications: Hematology/Vascular, Genitourinary, Neurologic, Outcome, and System Issue. Each column contains a list of specific complications, such as Hemolytic Anemia (Drug, etc), Acute Tubular Necrosis, Pneumocephalus, Euglycemic DKA, and Catheter Related Complication. The interface is designed for easy navigation and data entry.

Attending Evaluation: Resident Critique

Surgeon: [Redacted] Currently Selected Complications: [Redacted]
 DOB: [Redacted] RmTime: [Redacted]
 AGE: [Redacted] SrgTime: [Redacted]
 MRN: [Redacted] DOS: [Redacted] SEX: [Redacted] CaseType: [Redacted]
 Procedure: [Redacted]

Buttons: Delete Selected, Email Resident Grade, Save And Back, Cancel

System Complications 2

Cardiovascular	Pulmonary	Infection	Gastrointestinal	MusculoSkeletal	Head and Neck
Cardiac Ischemia	Hydropneumothorax	Variceal Bleeding	Chyle Leak Abdomen	Contracture	Patient Complaint
Pneumomediastinum	Negative Pressure _ PostObstruc tive Pulm Edema	Abscess, Intrathoracic	PO intolerance	Delayed Union	Emotional Psych Trauma
Angina	Hypospemia	Arthritis, Non-Septic	Esophagitis	Failure of Hardware	Acute Respiratory Distress
Arrhythmia	Air Leak prolonged	Bacteremia	Obnuzator (food impaction) obstr uction	Loss of Fixation	Anosmia (CN I)
Bradycardia	Arterial Ischemia Central	C.Difficile abx Colitis	Staple Line Leak	Loss Range of Motion	Atlanto-axial Instability (torticol lis, Grisel syndrome)
Cardiogenic Shock	Arterial Ischemia Peripheral	Cellulitis	Parastomal Hernia	Malunion	Carotid Blowout
CHF LVD Congestive Heart Fail ure	Aspiration	CLABSI	Abdominal Compartment Syndr ome	Non-Union	CN V Palsy (Trigeminal Neurop athy)
Dehydration	Asthma	Covid 19-Hospital Acquired	Anastomotic Dehiscence	Osteonecrosis	CN IV palsy (Trochlear)
Fluid Imbalance	Atelectasis	Dehiscence Wound	Anastomotic bleed_hemorrhage	Prosthesis Infection	CN IX Injury_Palsy (Glossophar yngeal)
Hypertension	BronchoPulmonary Fistula	Dehiscence/Evisceration	Anastomotic Leak	Prosthesis Loosening	CN IX Injury_Palsy (Glossophar yngeal)
Hypotension	Chyle Leak Neck	Explant Implant	Anastomotic Stricture	Prosthesis Malposition	CN XI Injury_Palsy(Spinal Acc essory)
hypovolemic shock	Chylothorax	Fever, prolonged	Ascites	Prosthesis Dislocation	CN XII Injury_Palsv (Hypooglos)
Lymphedema	COPD Exacerbation	Free Flap Failure		Prosthetic(Peri) Fracture	
		MSOF		Septic Joint	

Figure 1. Click-based surgical complication reporting platform.

Attending Evaluation: Resident Critique

Surgeon: [Redacted] Currently Selected Complications: [Redacted]
 DOB: [Redacted] RmTime: [Redacted]
 AGE: [Redacted] SrgTime: [Redacted]
 MRN: [Redacted] DOS: [Redacted] SEX: [Redacted] CaseType: [Redacted]
 Procedure: [Redacted]

Buttons: Delete Selected, Email Resident Grade, Save And Back, Cancel

Complications | **Attending Evaluation**

AttendingWriteUp | Case Analysis | Clavien Dindo Score | Harm Scale | Cognitive Bias (A-O) | Cognitive Bias (P-Z) | Management | Core Competencies | AttendingEvalSummary

Patient Centered Care	Medical Management	System Based Practice
Intra-Operative Patient Care: Performance of Procedures	Anatomy	Patient Safety and Quality Improvement (QI)
Intra-Operative Patient Care: Technical Skills	Pathophysiology and Treatment	Surgeon Role in Health Care Systems
Patient Evaluation and Decision Making		System Navigation for Patient-Centered Care
Post-Operative Patient Care		
Practice Management Improvement	Professionalism	Interpersonal/Communication Skills
Evidence-Based and Informed Practice	Accountability/Conscientiousness	Interprofessional and Team Communication
Reflective Practice and Commitment to Personal Growth	Self-Awareness and Help-Seeking	Communication within Health Care Systems
	Ethical Principles	Patient- and Family-Centered Communication
	Administrative Tasks	

Enter Write-Up Case Analysis / Critique | ClavienDindoScale | Harm Scale | Cognitive Bias (A - O) | Cognitive Bias (P - Z) | Management | Core Competencies

Responsible Party	Timing	Diagnosis	Technique	Judgment
Physician Team System Services Patient Compliance Patient Disease	Pre-Op Intra-Op Post-op Outpatient	Error Diagnosis Delay Diagnosis Inappropriate Test Failure to Act on Test Inadequate/Incomplete Pre-Op W/U	Error Technique/Performance Error Choice Operation Inappropriate Technique Error Tissue Handling/Tissue Injury Error Anatomy/Recognition	Error Judgment Inappropriate Indication for Surgery Inappropriate Patient Selection Error Timing Intervention Error Assessment Pathophysiology
Communication	Therapy/Prophylaxis	Professionalism	System Issues	
With Family/Informed Consent/Plan of Care Error in Signout/Handoff With Team Members With Consultants End of Life/Palliative Care Issues	Omission of Care Failure Prophylaxis Inadequate Monitoring Inadequate FollowUp Pre or Post Procedure Care	Acknowledged/Need for Help Identified/Accepted Responsibility Work Hours/Staffing issue Supervision Issue Credentialling Issue	Equipment Failure/Supplies Medication Issue (Drug, Dose, Patient, Route) Near Miss Pre or Post Procedure Care Complication Another Service/Other Hospital Nursing Issue Anesthesia Complication	

Figure 2. Click-based error reporting platform.

Thirty-seven subtypes of error are specified. Data was electronically collected on cases with complications and reviewed at weekly validation meetings by a committee of attendings (including the site Chief of Surgery, Program Director, divisional leadership, senior and junior residents, nursing and quality). The process was prospective and comprehensive. Cases were presented by residents responsible for care and data was validated by the above committee. Data was evaluated as the incidence of error across the seven domains, thirty-seven subtypes, and for specific procedures: bariatrics, cholecystectomy, colectomy, hepatic procedures, hiatal hernia, ileostomy closure, pancreatectomy and ventral hernia/component cases.

RESULTS

The study population included 25,857 general surgery/vascular cases, 4,005 complications, 1,473 cases with complications, and 3,268 errors or 2.2 errors per complication case. The number and incidence

of error by domain category was technical (831, 25.4%), therapy/prophylaxis (598, 18.3%), judgment (509, 15.6%), diagnosis (378, 11.6%), system issues (379, 11.3%), communication (356, 10.9%), and professionalism (226, 6.9%) (Table 1).

The most common subtypes for each domain were 'error technical performance' (54.2%), 'post procedure care MD Team' (79.4%), 'error medical judgment' (50.5%), 'delay diagnosis' (11.6%), 'pre/post procedure care Nursing Team' (59.7%), 'error in handoff team members' (10.9%), and 'identified/accepted responsibility' (62%), respectively.

All of the operative procedures evaluated mimicked the error profile of the domain and subtype categories with some differences in emphasis. We calculated the proportion of operations with complications that had errors. The median sub-specialty procedures had a rate of 0.83, with a range of only 0.77 to 0.89. Thus, a large proportion of operations with complications had at least one error, and this did not vary much across sub-specialty procedure

Table 1. Breakdown of errors by number and percent of cases.

Error Category	Number of Cases with Reported Error	Percent of Cases with Reported Error
Technique	831	25.43%
Error Technique/Performance	450	54.15%
Error Tissue Handling/Tissue Injury	189	22.74%
Error Anatomy/Recognition	67	8.06%
Inappropriate Technique	64	7.70%
Error Choice Operation	61	7.34%
Therapy/Prophylaxis	598	18.30%
Pre or Post Procedure Care (MD Team)	475	79.43%
Inadequate Monitoring	38	6.35%
Omission of Care	31	5.18%
Inadequate FollowUp	29	4.85%
Failure Prophylaxis/Failure to Follow Checklist	25	4.18%
Judgement	509	15.58%
Error Medical Judgment	257	50.49%
Error Timing Intervention	114	22.40%
Error Assessment Pathophysiology	83	16.31%
Inappropriate Patient Selection	43	8.45%
Inappropriate Indication for Surgery	12	2.36%
Diagnosis	378	11.57%
Delay Diagnosis	209	55.29%
Failure to Act on Test	64	16.93%
Error Diagnosis	50	13.23%
Inadequate/Incomplete Pre-Op W/U	41	10.85%

Error Category	Number of Cases with Reported Error	Percent of Cases with Reported Error
Inappropriate Test	14	3.70%
System Issues	370	11.32%
Pre or Post Procedure Care (Nursing Team)	221	59.73%
Complication Another Service/Other Hospital	66	17.84%
Medication Issue (Drug, Dose, Patient, Route)	32	8.65%
Anesthesia Complication	18	4.86%
Equipment Failure/Supplies	11	2.97%
Near Miss	10	2.70%
Nursing Error	10	2.70%
Operating Room Delay	2	0.54%
Communication	356	10.89%
Error Handoff Team Members	144	40.45%
With Family/Informed Consent/Plan of Care	105	29.49%
Error Handoff Signout with Consultants	63	17.70%
End of Life/Palliative Care Issues	30	8.43%
Error in Signout/Handoff	14	3.93%
Professionalism	226	6.92%
Identified/Accepted Responsibility	140	61.95%
Acknowledged/Need for Help	77	34.07%
Supervision Issue	6	2.65%
Work Hours/Staffing issue	2	0.88%
Credentiailling Issue	1	0.44%

DISCUSSION

Medical error, as an inherent part of healthcare, continues to be a significant challenge in clinical practice, with profound implications for patient safety, quality of care, and healthcare outcomes. Our study offers critical insights into the types and distribution of errors in surgical practice, supporting the idea that errors leading to patient harm are rarely the result of a single factor. Rather, they arise from a complex interplay of human, systemic, and procedural elements. In addition, system issues accounted for only 11% of errors, highlighting that provider-based technical and cognitive errors accounted for nearly 90% of errors. These findings expand upon existing literature, which has similarly emphasized the multifactorial nature of medical error and its relationship to various contributing factors, including clinical judgment, technical performance, communication breakdowns, and system-level issues.

The most common categories of errors observed in our study were technical and judgment errors, followed by errors in therapy, communication, and system failures. Specifically, technical performance and post-procedure care errors were the most frequently identified, which aligns with prior studies identifying the importance

of technical proficiency and follow-up care in mitigating patient harm. Furthermore, the relationship between different surgical specialties and error profiles revealed that while specific error types may be more common in certain procedures (e.g., technical errors in complex surgeries such as pancreatectomy), the overall frequency of errors across different domains remained relatively consistent. This suggests that errors are a common feature across surgical disciplines, and strategies to mitigate them should be broadly applicable across various surgical specialties.

One of the most striking observations in this study is the high incidence of errors even in cases that did not result in harm. This highlights an important distinction in error research: not all errors lead to adverse outcomes, but the potential for harm remains ever-present. It underscores the need for a broader approach to error prevention, one that encompasses not just harm reduction but also efforts to minimize the frequency of error occurrence overall. While some errors may be benign, their presence signals opportunities for improvement in clinical practice, communication, and system design.

Our data also underscore the importance of standardized error reporting and classification systems. The template and error

domains used in this study allowed for a systematic approach to error identification and analysis, facilitating the detection of recurring error patterns and trends. Although data is not specifically presented in this paper, the error data collected was automatically applied to individual procedure classes and to individual surgeons in the form of periodic reviewable report cards. Such a framework can serve as a valuable tool for quality improvement initiatives, enabling healthcare institutions to identify areas for targeted interventions and develop evidence-based strategies to reduce error rates.

CONCLUSION

These data suggest that patterns of error can be accurately described utilizing a standardized error profile template to analyze post-operative complications. Operation types with more errors had more complications. By understanding the distribution of error types, providers and colleagues may be informed of their individual propensity for error, and systems can be more focused in quality improvement initiatives based on patterns specific to the case type. In addition, system issues accounted for only 11% of errors, highlighting that provider-based technical and cognitive errors accounted for nearly 90% of errors.

The use of standardized frameworks, as demonstrated in this study, can aid in the development of targeted interventions and ultimately improve the safety and quality of care provided to patients. Moving forward, healthcare institutions should prioritize error reduction strategies, focusing on education, cognitive performance, prevention and the optimization of clinical workflows, communication, and team dynamics to reduce the frequency of errors and improve patient outcomes in surgical settings.

Furthermore, the study emphasizes the need for robust error reporting systems that can identify patterns of mistakes across different error domains, providing valuable data to inform continuous quality improvement efforts. By recognizing the prevalence of medical error, embracing a culture of transparency, and fostering collaborative approaches to patient care, healthcare providers can take important steps toward reducing preventable harm in surgical practice.

DISCLOSURES

Antonacci: President of Outcomes Management Systems, LLC

ACKNOWLEDGEMENT

None

FUNDING

The author received no financial support for the research, authorship, and/or publication of this article.

CONFLICT OF INTEREST

The author declares no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

REFERENCES

1. Makary MA, Daniel M. Medical error-the third leading cause of death in the US. *BMJ*. 2016;353:i2139.
2. Institute of Medicine (US) Committee on Quality of Health Care in America. *To Err is Human: Building a Safer Health System*. Kohn LT, Corrigan JM, Donaldson MS, editors. Washington (DC): National Academies Press (US); 2000.
3. Marsh KM, Turrentine FE, Jin R, et al. Judgment Errors in Surgical Care. *J Am Coll Surg*. 2024;238(5):874-879.
4. Marsh KM, Turrentine FE, Knight K, et al. Defining and Studying Errors in Surgical Care: A Systematic Review. *Ann Surg*. 2022;275(6):1067-1073.
5. Antonacci AC, Lam S, Lavarias V, et al. A morbidity and mortality conference-based classification system for adverse events: surgical outcome analysis: part I. *J Surg Res*. 2008;147(2):172-7.
6. Xu J, Sherry ML, Kochaneck KD, et al. Deaths: final data for 2013. *National Vital Statistics Reports*. 2016;64(2).
7. Antonacci AC, Lam S, Lavarias V, Homel P, Eavey RA. A report card system using error profile analysis and concurrent morbidity and mortality review: surgical outcome analysis, part II. *J Surg Res*. 2009;153(1):95-104.
8. Antonacci AC, Dechario SP, Antonacci C, et al. Cognitive Bias Impact on Management of Postoperative Complications, Medical Error, and Standard of Care. *J Surg Res*. 2021;258:47-53.
9. Antonacci AC, Dechario SP, Rindskopf D, Husk G, Jarrett M. Cognitive bias and severity of harm following surgery: Plan for workflow debiasing strategy. *The American Journal of Surgery*. 2021;222(6):1172-7.
10. Carsky K, Rindskopf D, Patel VM, et al. Using Concurrent Complication Reporting to Evaluate Resident Critical Thinking and Enhance Adult Learning. *J Surg Educ*. 2024;81(5):702-712.
11. Antonacci AC, Farrell A, Portelli K, et al. Optimizing Complication Self-Reporting Methodologies Improves Standard of Care and Quality. *J Surg Res*. 2025;309:277-287.